

pub C4

- (b) reacting an admixture of the oxazolidinone derivative and L-phenylalanine or L-phenylalanine methyl ester in a second solvent for a time and at a temperature sufficient to produce N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester.

[illegible]

- SubH
A3

4. The process according to claim 1, wherein the first solvent is selected from the group consisting of tetrahydrofuran, diethyl ether, t-butyl methyl ether, ethyl acetate, dioxane, toluene, butyl acetate, methyl

acetate, dichloromethane, dimethylformamide, dimethylsulfoxide and combinations thereof.

5. The process according to claim 1, wherein the ratio of N-(3,3-dimethylbutyl)-L-aspartic acid to the carbonyl compound is from about 1:1 to about 1:4.

6. The process according to claim 1, wherein the temperature sufficient to produce the oxazolidinone derivative is from about 20°C to about 150°C.

7. The process according to claim 6, wherein the temperature sufficient to produce the oxazolidinone derivative is from about 22°C to about 70°C.

8. The process according to claim 1, wherein the time sufficient to produce the oxazolidinone derivative is from about 1 hour to about 48 hours.

9. The process according to claim 8, wherein the time sufficient to produce the oxazolidinone derivative is from about 12 hours to about 24 hours.

10. The process according to claim 1, wherein the admixture of N-(3,3-dimethylbutyl)-L-aspartic acid and a carbonyl compound or an activated carbonyl compound further comprises a catalyst.

~~11. The process according to claim 10, wherein the catalyst is p-toluenesulfonate.~~

~~12. The process according to claim 1, wherein the admixture of N-(3,3-dimethylbutyl)-L-aspartic acid and a carbonyl compound or an activated carbonyl further comprises an acid.~~

Subl
A5
13. The process according to claim 12, wherein the acid is selected from the group consisting of formic acid, acetic acid, p-toluene sulfonic acid, methane sulfonic acid and combinations thereof.

14. The process according to claim 1, wherein the second solvent is selected from the group consisting of tetrahydrofuran, diethyl ether, t-butyl methyl ether, ethyl acetate, dioxane, toluene, butyl acetate, methyl acetate, dichloromethane, dimethylformamide, dimethylsulfoxide and combinations thereof.

15. The process according to claim 1, wherein the first solvent and the second solvent are the same.

16. The process according to claim 1, wherein the ratio of L-phenylalanine or L-phenylalanine methyl ester to the oxazolidinone derivative is from about 1:1 to about 1:2.

17. The process according to claim 1, wherein the temperature sufficient to produce N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is from about 0°C to about 50°C.

18. The process according to claim 17, wherein the temperature sufficient to produce N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is from about 22°C to about 40°C.

19. The process according to claim 1, wherein the time sufficient to produce N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is from about 1 hour to about 48 hours.

20. The process according to claim 19, wherein the time sufficient to produce N-[N-(3,3-dimethylbutyl)-L-

COPIED FROM THE FILE

~~α-aspartyl]-L-phenylalanine 1-methyl ester is from about 12 hours to about 24 hours.~~

Parameter	Estimate	Standard Error	z	P	95% CI
Intercept	0.000	0.000	0.000	1.000	0.000, 0.000
Age	0.000	0.000	0.000	1.000	0.000, 0.000
Gender	0.000	0.000	0.000	1.000	0.000, 0.000
Education	0.000	0.000	0.000	1.000	0.000, 0.000
Income	0.000	0.000	0.000	1.000	0.000, 0.000
Health	0.000	0.000	0.000	1.000	0.000, 0.000
Marital	0.000	0.000	0.000	1.000	0.000, 0.000
Religion	0.000	0.000	0.000	1.000	0.000, 0.000
Occupation	0.000	0.000	0.000	1.000	0.000, 0.000
City	0.000	0.000	0.000	1.000	0.000, 0.000
State	0.000	0.000	0.000	1.000	0.000, 0.000
Country	0.000	0.000	0.000	1.000	0.000, 0.000
Year	0.000	0.000	0.000	1.000	0.000, 0.000
Constant	0.000	0.000	0.000	1.000	0.000, 0.000